

Dear Physician:

Please prescribe Twinject™ auto-injectable epinephrine 1:1000 to our Wilderness Medicine Outfitters training graduate, _____, who has completed a course in the recognition and treatment of anaphylaxis, as well in the use of injectable medications. The graduate to whom you will be prescribing and/or delegating the use of Twinject™ to treat patients who are experiencing a life-threatening allergic reaction in a remote or wilderness setting has been specifically trained to use this device. Please do not substitute EpiPen for Twinject™, as a single EpiPen contains only one dose of epinephrine, whereas each Twinject™ device contains two doses of epinephrine in a durable and compact carrying case. Approximately thirty percent of anaphylaxis episodes will require treatment with more than one dose of epinephrine. The first dose of epinephrine in Twinject™ should be used as early as possible after the onset of signs and/or symptoms of anaphylaxis. If the patient is not improved or is worse about five minutes after the first dose has been administered, or if new signs and/or symptoms develop, the second dose of epinephrine contained in Twinject™ should be given. This is especially important with respiratory and/or cardiovascular system involvement. The graduate has been trained on the importance of evacuation for evaluation in a medical facility when anaphylaxis has occurred.

Please call with any questions.

**Sincerely,
Carl Weil
Director Wilderness Medicine Outfitters
303-688- 5176**